



AGCHIMP™

COVERS UP TO 5,000 Sq. Ft.

18-0-9

30% Slow Release with 2% Iron

PRODIAMINE GROUP 3 HERBICIDE

PRODIAMINE 0.29% PLUS TURF FERTILIZER

- FOR PREEMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF (excluding golf course putting greens)
STOPS MANY WEED GRASSES BEFORE THEY GROW
PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL

Table with 2 columns: Ingredient and Percentage. Prodiamine* 0.29%, Other Ingredients 99.71%, Total 100.00%. Includes CAS # 29091-21-2, EPA Reg No. 53883-170-102434, EPA Est. No. 102434-GA-1.

CAUTION

KEEP OUT OF REACH OF CHILDREN
Read the entire label carefully before using this product.

Table with 2 columns: Situation and First Aid instructions. Includes sections for IF IN EYES, IF ON SKIN OR CLOTHING, IF SWALLOWED, IF INHALED, and HOT LINE NUMBER.

See the Precautionary Statements, and Hazards to Humans and Domestic Animals for additional information.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow this product to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather will help to ensure that wind does not blow this product off the treated area.

Table with 2 columns: Nutrient and Percentage. 18-0-9 GUARANTEED ANALYSIS. Total Nitrogen (N) 18.00%, Urea Nitrogen* 18.00%, Total Soluble Potash (K2O) 9.00%, Chlorine (Cl) MAX 6.75%, Iron (Fe) 2.00%, Water Soluble Iron 0.16%.

DERIVED FROM: Urea, Polymer Coated Urea, Potassium Sulfate, Iron EDTA
*4.86% Slowly Available Nitrogen from Polymer Coated Urea.

Distributed & Guaranteed By: AgChimp LLC, 127 Riverside Dr SW, Cartersville GA 30120, 770-334-9442



NET WEIGHT 15 lbs. (6.80kg)

CONDITIONS OF SALE AND WARRANTY: The directions for use of this product are based on the conditions of sale and warranty. The user assumes all responsibility for the use of this product. Copy, reprint, or other unauthorized use is prohibited without the written consent of AgChimp LLC or the seller.

Table with 2 columns: Turf Grass Species and Minimum Annual Rate (lb/1,000 sq. ft.). Includes Creeping Bentgrass, Creeping Red Fescue, Kentucky Bluegrass, etc.

USE RATE: Months Before Overseeding South. Includes instructions for application rates and timing.

WHEN TO OVERSEED AFTER APPLICATION: This product will inhibit turf germination if applied too soon after application. Follow rates and intervals in table for best results.

APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been calibrated to apply 2.25 to 3.50 pounds/acre. Followed after 60 to 90 days by a second application of 5.2 to 7.9 pounds/acre.

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds. It will not control established weeds.

MINIMUM AND MAXIMUM APPLICATION RATES (By Turf Grass Species): Table with 2 columns: Species and Rate (lb/1,000 sq. ft.).

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per acre is given for each turf grass species. Do not exceed the maximum yearly rate.

USE RESTRICTIONS: Do not apply to newly seeded turf within 60 days after seeding or until after the second mowing. Do not apply to putting greens or areas where divots are likely to be taken.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may result in turf damage if applied in excess.



Table with 2 columns: Spreader Settings and Rate (lb/1,000 sq. ft.). Includes settings for various spreader models like LESCO, Craftsman, etc.

APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been calibrated to apply 2.25 to 3.50 pounds/acre. Followed after 60 to 90 days by a second application of 5.2 to 7.9 pounds/acre.

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds. It will not control established weeds.

MINIMUM AND MAXIMUM APPLICATION RATES (By Turf Grass Species): Table with 2 columns: Species and Rate (lb/1,000 sq. ft.).

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per acre is given for each turf grass species. Do not exceed the maximum yearly rate.

USE RESTRICTIONS: Do not apply to newly seeded turf within 60 days after seeding or until after the second mowing. Do not apply to putting greens or areas where divots are likely to be taken.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may result in turf damage if applied in excess.

APPLICATION DIRECTIONS: Apply this product uniformly, using a suitable spreader that has been calibrated to apply 2.25 to 3.50 pounds/acre. Followed after 60 to 90 days by a second application of 5.2 to 7.9 pounds/acre.

WHEN TO APPLY TO ESTABLISHED TURF: This product must be applied prior to germination of weed seeds. It will not control established weeds.

MINIMUM AND MAXIMUM APPLICATION RATES (By Turf Grass Species): Table with 2 columns: Species and Rate (lb/1,000 sq. ft.).

RATES OF APPLICATION: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per acre is given for each turf grass species. Do not exceed the maximum yearly rate.

USE RESTRICTIONS: Do not apply to newly seeded turf within 60 days after seeding or until after the second mowing. Do not apply to putting greens or areas where divots are likely to be taken.

USE PRECAUTIONS: The following precautions apply to the use of this product in turf grasses and lawns. Application of this product may result in turf damage if applied in excess.