



Bermudagrass, also known as couch grass in Australia and New Zealand, is a grass found worldwide. Classified as a warm season grass, bermudagrass is typically active growing from mid-March through last frost. Following last frost, bermudagrass transitions into its dormant state which will appear with a straw looking appearance. Bermudagrass will transition from its dormant state when soil temperatures begin to warm in the spring.

Mowing Height: 1-2 inches.

Water Requirements: 1" / week

Fertilization: 2-5 lbs/Nitrogen/Year

Ideal Soil pH: 5.5-6.5



AgChimp™ Basic Maintenance Program	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
5-0-30 2% Iron .29% Prodiamine												
18-0-6 30% Slow Release .29% Prodiamine												
24-0-11 30% Slow Release 2% Iron												
AgChimp™ Sprayable Iron												
AgChimp™ pHix Sprayable Limestone												
AgChimp™ PunchCa												
AgChimp™ Three-D Selective Herbicide												
AgChimp™ Conduit Selective Herbicide												
As Needed for Broadleaf Weed Control												
As Needed for Grassy Weed Control												

AgChimp™ PLUS Maintenance Program	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
5-0-30 2% Iron .29% Prodiamine												
18-0-6 30% Slow Release .29% Prodiamine												
24-0-11 30% Slow Release 2% Iron												
AgChimp Sprayable Iron												
AgChimp pHix Sprayable Limestone												
AgChimp PunchCa												
AgChimp JuiceRx												
AgChimp Strobe+ G												
5-0-15 2% Iron .069% Bifenthrin Insecticide												
AgChimp™ Three-D Selective Herbicide												
AgChimp™ Conduit Selective Herbicide												
As Needed for Broadleaf Weed Control												
As Needed for Grassy Weed Control												

